

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

1-23-2006

Soybean aphids and K-deficient soybeans: Why are there aphids in my field and not my neighbor's?

Matthew E. O'Neal

Iowa State University, oneal@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

O'Neal, Matthew E., "Soybean aphids and K-deficient soybeans: Why are there aphids in my field and not my neighbor's?" (2006). *Integrated Crop Management News*. Paper 1182.
<http://lib.dr.iastate.edu/cropnews/1182>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

INTEGRATED CROP MANAGEMENT

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2006 > IC-496 (1) -- January 23, 2006

Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

[Image Gallery](#)

Printable Version

Printable version of this page

Related Articles

How do uncertain prices influence phosphorus and potassium fertilization this fall?
October 1, 2007

Aphids attack!
August 6, 2007

Extension agronomists update soybean aphid

Soybean aphids and K-deficient soybeans: Why are there aphids in my field and not my neighbor's?

by Matt O'Neal, Department of Entomology

There were frequent reports of fields infested above 250 soybean aphids per plant adjacent to those with lower populations that never reached threshold. The lowest aphid populations were seen in the southern tier of Iowa counties. However, Mark Carlton, extension field crop specialist in southeast Iowa, called me several times during the season to report that he had scouted fields that were above threshold. By the third call, it became a bit of a routine, with each field sharing some remarkable similarities. All these soybean fields were planted in either alfalfa or sod in 2004. This is interesting since soybeans following alfalfa often suffer from potassium deficiency. And to complete the puzzle, recent research conducted at the University of Wisconsin and Michigan State University indicates that soybeans grown in potassium-deficient soils may be more at risk for soybean aphid outbreaks. What follows is a brief explanation of why soybean aphids reproduce faster on K-deficient soybeans.

But before I get to the K-deficient soybeans, let's consider the feeding habits of soybean aphids. This species is like most aphids: they get all of their nutrition from one source--the phloem of a single species of plant. To put this in some perspective, their meat, potatoes, and dessert are all taken from one source. Since phloem is low in vital nutrients, like nitrogen, it is thought that the aphids make up for this apparently unbalanced diet by ingesting a lot fluid to get all the nutrients they need to grow and reproduce. Any excess fluid and sugar may contribute to the honeydew that aphids shed while feeding on plants.

Given the low concentrations of nitrogen in the phloem of most plants, changes in its concentration can have dramatic changes in aphid growth and reproduction. For most aphid/plant interactions, the greatest amount of nitrogen in phloem corresponds to two periods in a plant's life--vegetative growth and senescence. These are times when nitrogen is flowing in the plant, moving to growing leaves or shuttled from senescing leaves. These are also times when aphids tend to grow and reproduce in the greatest number. Between these periods of robust growth, the amount of nitrogen available in the phloem is not as great. This may explain why soybean aphids look "anemic" or "whitish" in late July and early August since soybean vegetative growth is completed and the availability of nitrogen in the phloem may be limited.

Another factor that can influence nitrogen availability is stress. One type of stress that has been shown to favor aphid growth is K-deficiency. Recently, a group of entomologists at

situation

July 30, 2007

Newly revised aphids
publication available

July 23, 2007

Mid-July soybean aphid
update

July 16, 2007

Soybean aphids exceed
the economic threshold
in northeast Iowa

July 9, 2007

Monitor soybean aphid
populations on PIPE

July 2, 2007

Soybean aphid numbers
increase...and decrease

July 2, 2007

Early soybean aphid
outbreak--to the east

June 11, 2007

Soybean aphids found in
Minnesota and
Wisconsin

June 4, 2007

PIPE: Pest Information
Platform for Extension
and Education

March 26, 2007

the University of Wisconsin showed that aphids on K-deficient soybeans produce 39 percent more nymphs than on healthy plants. In a separate study, Chris DiFonzo and her graduate student Abigail Summers at Michigan State University have shown that the amino acid profile in K-deficient soybeans contains a greater amount of amino acids that aphids need for optimal growth. So it appears that although a K-deficient plant is stressed, it is a better host for soybean aphids than a healthy plant.

To what extent this contributes to local differences in soybean aphid populations is not yet clear. There are potentially several plant-stress factors that could lead to similar changes in soybean health and subsequent aphid population growth. For example, Greg Tylka, extension nematologist and plant pathologist; Felicitas Avendano, researcher; and I are collaborating to determine if soybean cyst nematodes also stress soybeans in such a way as to exacerbate aphid population growth. We hope to identify factors that would increase the likelihood of soybean aphid outbreaks and help growers better predict what fields may be at risk for economic populations.

This article originally appeared on page 6 of the IC-496 (1) -- January 23, 2006 issue.

Updated 11/08/2006 - 5:35pm